## 64548

# Dilithic Breccia 8.49 grams



Figure 1: Photo of 64548. Mm scale. S72-55353.

#### Introduction

64548 is a rake sample (walnut) from the boulder field at station 4 on Stone Mountain – see section on 64501. It is not known if it is local, or if it is part of the ejecta that landed there. It is a fine grain mix of chalky white anorthite and dark aphanitic impact melt (figure 1).

## **Petrography**

Warner et al. (1973) classify 64548 as a "light matrix breccia". However, McKinley et al. (1983) and others realized that 64548 was essentially the same as the sequence 64535, 64546, because of the distinctive light and dark texture and clean boundary between the two lithologies (figure 2).

Ryder and Norman (1980) reported that the matrix was crushed anorthositic material intimately mixed with patches of dark impact melt.

### **Chemistry**

McKinley et al. (1983) and Floran et al. (1976) reported analyses of 64548 (table 1). Ryder and Norman (1980)


referred to unpublished analyses by Blanchard (figure 3).

## **Other Studies**

Pearce and Simonds (1974) determined the magnetic properties of 64548 and Gooley et al. (1973) determined the Ni and Co of iron grains and schreibersite.

## **Processing**

There is only one thin section of 64548.






Figure 2: Photomicrograph of thin section 64548,4. Width of field is 2 mm.




Figure 3: Normalized rare-earth-element diagram for 64548 (Blanchard).

Table 1. Chemical composition of 64548.

reference weight	Floran76		Blanchard		McKinley83	
SiO2 % TiO2 Al2O3 FeO MnO MgO CaO Na2O K2O P2O5 S % sum	45.28 0.43 27.67 4.47 5.67 15.79 0.46 0.13	(c) (c) (c) (c) (c) (c)			47.6 0.88 21.24 5.9 0.09 10.27 12.83 0.52 0.27	(a) (a) (a) (a) (a) (a) (a) (a)
(a) Sc ppm V Cr			6.78	(b)		
Co Ni Cu Zn Ga Ge ppb As Se Rb Sr Y Zr Nb Mo Ru Rh Pd ppb Ag ppb Cd ppb In ppb Sn ppb Sb ppb Te ppb			24.5 380	(b) (b)		
Cs ppm Ba La Ce Pr Nd Sm Eu Gd Tb Dy Ho Er Tm			14.6	(b)		
Yb Lu Hf Ta W ppb Re ppb Os ppb Ir ppb Pt ppb Au ppb Th ppm U ppm			0.67	(b)		
technique:	(a) broa	ad be	am e p	robe.	(b) INAA	. (c)

#### **References for 64548**

Butler P. (1972a) Lunar Sample Information Catalog Apollo 16. Lunar Receiving Laboratory. MSC 03210 Curator's Catalog. pp. 370.

Gooley R.C., Brett R. and Warner J.L. (1973) Crystallization history of metal particles in Apollo 16 rake samples. *Proc.* 4<sup>th</sup> Lunar Sci. Conf. 799-810.

Korotev R.L. (1994) Compositional variation in Apollo 16 impact melt breccias and inferences for the geology and bombardment history of the central highlands of the Moon. *Geochim. Cosmochim. Acta* **58**, 3931-3969.

LSPET (1973b) The Apollo 16 lunar samples: Petrographic and chemical description. *Science* **179**, 23-34.

LSPET (1972c) Preliminary examination of lunar samples. *In* Apollo 16 Preliminary Science Report. NASA SP-315, 7-1—7-58.

McKinley J.P., Taylor G.J., Keil K., Ma M.-S. and Schmitt R.A. (1984) Apollo 16: Impact sheets, contrasting nature of the Cayley Plains and Descartes Mountains, and geologic history. *Proc.* 14<sup>th</sup> Lunar Planet. Sci. Conf. in J. Geophys. Res. **89**, B513-B524.

Pearce G.W. and Simonds C.H. (1974) Magnetic properties of Apollo 16 samples and implications for their mode of formation. *J. Geophys. Res.* **79**, 2953-2959.

Phinney W. and Lofgren G. (1973) Description, classification and inventory of Apollo 16 rake samples from stations 1, 4 and 13. Curators Office.

Ryder G. and Norman M.D. (1980) Catalog of Apollo 16 rocks (3 vol.). Curator's Office pub. #52, JSC #16904

Sutton R.L. (1981) Documentation of Apollo 16 samples. In Geology of the Apollo 16 area, central lunar highlands. (Ulrich et al. ) U.S.G.S. Prof. Paper 1048.

Warner J.L., Simonds C.H. and Phinney W.C. (1973b) Apollo 16 rocks: Classification and petrogenetic model. *Proc.* 4<sup>th</sup> *Lunar Sci. Conf.* 481-504.

technique: (a) broad beam e probe, (b) INAA, (c) fused bead e probe.